



HOW TO USE

DO NOT exceed 3m between bait stations for mice. DO NOT place bait stations in the open.

SITUATION	PEST	RATE	HOW TO APPLY
In and around domestic homes, animal houses, farms and public service buildings.	Mice (<i>Mus musculus</i>)	Place one bait station per infested area	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least two weeks is necessary to reduce mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

HOW TO PREPARE

Determine areas where mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where mice or their signs have been observed. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce mouse numbers.

CAUTION

DO NOT use product in areas accessible to children.
DO NOT allow bait to contaminate foodstuffs or food intended for human or animal consumption.
DO NOT apply bait to crops.
DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in locked room or place away from children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specially marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty box and product should not be burnt. Dispose of carcasses safely by burying and burning in an approved landfill.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear rubber gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126).

MSDS

For further information please refer to the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

The use of TOMCAT ALL-WEATHER BLOX is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Competition and Consumer Act 2010.

073015

APVMA APPROVAL NO: 49776/58197